

- GLEASON, H. A. 1952. The New Britton and Brown Illustrated Flora of Northeastern United States and Adjacent Canada. 2: 536–537.
- JACKSON, B. D. 1912. Index to the Linnaean Herbarium. Linnaean Society, London.
- KELLER, R. 1925. *Ascyrum* in Engler and Prantl, Die Natürlichen Pflanzenfamilien. 2 Auflage, 21 Band, 175.
- LANJOUW, J. and F. A. STAFLEU. 1956. The Herbaria of the World. Regnum Vegetabile. Vol. 6, ed. 3.
- LINNAEUS, C. 1737. Genera Plantarum. 231.
- . 1753. Species Plantarum. 2: 777–778.
- . 1763. Species Plantarum. ed. 2, 2: 1107–1108.
- MOHR, C. 1901. Plant Life of Alabama. Contr. U. S. Nat. Herb., 6.
- PLUMIER, C. 1703. Nova Plantarum Americanum Genera. 51.
- SAVAGE, S. 1945. A Catalog of the Linnaean Herbarium. Linnaean Society of London.
- SMALL, J. K. 1933. Manual of the southeastern Flora. 866–868.
- SPACH, E. 1836. Famille Hypericaceae. Histoire Naturelle des Végétaux. 5: 445–464.
- TORREY, J. and ASA GRAY. 1838–1840. Flora of North America. 1: 156–157, 671–672.
- VESTAL, P. A. 1937. The significance of comparative anatomy in establishing the relationship of the Hypericaceae to the Guttiferae and their allies. Philippine Journal of Science 64: 199–256.

HEDYOTIS MINIMA f. *ALBIFLORA*.—On April 6, 1955, when the author was collecting with Dr. Ronald L. McGregor, a colony of pure white flowered *H. minima* was discovered on shallow sandy soil, over sandstone, in a clearing of an upland scrub oak forest, in Woodson County, Kansas. The exact location was: Sec. 32, T25S, R15E. The specimens were scattered in a small area of this woodland.

Hedyotis minima (Beck) T. & G., forma ***albiflora***, f. nov.—Haec forma a planta typica speciei corollis albis recedit. This form differs from the typical form of the species in having white corollas. Type specimen *Lathrop and McGregor 35*, is in the Herbarium of the University of Kansas.—EARL LATHROP, DEPARTMENT OF BOTANY, UNIVERSITY OF KANSAS, LAWRENCE, KANSAS.